

2020 Reporting Portal Measures

Comagine Health provides the following quality and utilization measures to primary care medical groups and practices in Oregon via the secure online <u>Reporting Portal</u>. Some of the measure results (as noted below) are reported publicly on Comagine Health's <u>Compare Your Care</u> website. The data publicly reported in 2020 is from 2018; <u>Compare Your Care</u> will be updated with 2019 data in December 2020.

| Measure | Publicly Reported |
|-------------------------------------------------------------------------------------------------------------|----------------------|
| 30 Day All-Cause Readmissions, Unadjusted (age 18+) | |
| Acute Hospital Utilization – Medical (age 18+) | |
| Acute Hospital Utilization – Surgical (age 18+) | |
| Acute Hospital Utilization – Total (age 18+) | |
| Adolescent Well-Care Visits (age 12-21) | √ |
| Alcohol Misuse: Screening, Brief Intervention, Referral for Treatment (SBIRT) (age 12-17, age 12+, age 18+) | |
| Ambulatory care: Avoidable Emergency Department Visits, % of Total Visits (age 1-17, age 18+) | |
| Ambulatory care: Avoidable Emergency Department Visits, Rate per 1000 Patients (age 1-17, age 18+) | |
| Ambulatory care: Emergency Department Visits (age 0-17, age 18+) | |
| Ambulatory care: Outpatient Visits (age 0-17, age 18+) | |
| Annual Monitoring for Patients on Persistent Medications (age 18+) | |
| Appropriate Testing for Children with Pharyngitis (age 2-18) | ✓ |
| Appropriate Testing for Children with Upper Respiratory Infection (age 3 months to 18 years) | ✓ |
| Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis | ✓ |
| Breast Cancer Screening (age 50-74) | ✓ |
| Cervical Cancer Screening (age 21-64) | √ |
| Chlamydia Screening (age 16-20, age 21-24, age 16-24) | √ |
| Comprehensive Diabetes Care, Eye Exam Performed (age 18-75) | |
| Comprehensive Diabetes Care, HbA1c Testing (age 18-75) | ✓ |
| Comprehensive Diabetes Care, Medical Attention for Nephrology (age 18-75) | |

| Measure | Publicly Reported |
|-----------------------------------------------------------------------------------------------|----------------------|
| Developmental Screenings in the First 36 Months of Life (age 1, 2 and 3) | ✓ |
| Effective Contraceptive Use (age 15-17, age 18-50) | |
| Emergency Department Utilization (age 18+) | |
| Follow-up After Emergency Department Visit for Mental Illness (age 6+) | |
| Generic Prescription Fills: Antidepressants (age 18+) | ✓ |
| Generic Prescriptions Fills: Antihyperlipidemics (age 18+) | ✓ |
| Generic Prescription Fills: Antihypertensives (age 18+) | ✓ |
| Hospital Inpatient Admission Rates for Ambulatory Sensitive Conditions (PQI Measures) | |
| Medication Management for People with Asthma, Child (age 5-18) | ✓ |
| Statin Therapy for Patients with Cardiovascular Disease (males age 21-75, females age 40-75) | |
| Statin Therapy for Patients with Diabetes (age 40-75) | |
| Total Expenditures | |
| Total Inpatient Expenditures | |
| Total Specialty Expenditures | |
| Use of Imaging Studies for Low Back Pain (age 18-50) | ✓ |
| Use of Opioids at High Dosage in Persons Without Cancer (age 18+) | |
| Use of Opioids from Multiple Providers in Persons Without Cancer (age 18+) | |
| Use of Opioids at High Dosage and from Multiple Providers in Persons Without Cancer (age 18+) | |
| Well-Child Visits 0-15 Months, Five or More | |
| Well-Child Visits 0-15 Months, Six or More | √ |
| Well-Child Visits 3-6 Years | 1 |

Note: Comagine Health also publicly reports the Antidepressant Medication Management measure on a statewide basis but does not report the measure to medical groups or providers due to small denominator sizes.